

## Change of Variable

We will now discuss one last technique for solving non-linear first order differential equations, where the basic idea is to make a change of variables in order to recast the differential equation into one of the cases we have already solved. We'll begin with a special case of this method.

### 1. Homogeneous Equations of Degree Zero

The special form to be considered here is when the given differential equation

$$(11.1) \quad y' = f(x, y)$$

can be expressed as

$$(11.2) \quad y' = F\left(\frac{y}{x}\right) .$$

In such a case, we say the differential equation (11.1) is **homogeneous of degree zero**.

EXAMPLE 11.1.

$$(11.3) \quad y' = \frac{y^2 + 2xy}{x^2} = \left(\frac{y}{x}\right)^2 + 2\left(\frac{y}{x}\right)^{-1} = F\left(\frac{y}{x}\right) \text{ when } F(v) \equiv v^2 + \frac{2}{v}$$

EXAMPLE 11.2.

$$(11.4) \quad y' = \ln(x) - \ln(y) = -\ln\left(\frac{y}{x}\right) = F\left(\frac{y}{x}\right) \text{ when } F(v) \equiv -\ln(v)$$

EXAMPLE 11.3.

$$(11.5) \quad y' = \frac{x+y}{y-x} = \frac{1 + \left(\frac{y}{x}\right)}{\left(\frac{y}{x}\right) - 1} = F\left(\frac{y}{x}\right) \text{ when } F(v) \equiv \frac{1+v}{v-1}$$

To solve such equations we introduce a new variable which we will denote by  $v$ , to represent the ratio of  $y$  to  $x$ :

$$(11.6) \quad v = \frac{y}{x} .$$

We then have

$$(11.7) \quad \frac{dv}{dx} = -\frac{y}{x^2} + \frac{1}{x} \frac{dy}{dx}$$

or

$$(11.8) \quad \frac{dy}{dx} = x \left( \frac{y}{x^2} + \frac{dv}{dx} \right) = v + x \frac{dv}{dx} .$$

Hence equation (11.2) becomes

$$(11.9) \quad x \frac{dv}{dx} + v = F(v)$$

or

$$(11.10) \quad xdv = (F(v) - v) dx$$

or

$$(11.11) \quad \frac{dx}{x} = \frac{dv}{F(v) - v} \quad .$$

Integrating both sides of this equation yields

$$(11.12) \quad \ln|x| = \int \frac{dv}{F(v) - v} + C \quad .$$

Suppose now that the integral on the right hand side has been carried out so

$$(11.13) \quad H(v) = \int \frac{dv}{F(v) - v}$$

is some explicit function of  $v$ . Equation (11.12) becomes

$$(11.14) \quad \ln|x| - H(v) = C$$

or

$$(11.15) \quad \ln|x| - H\left(\frac{y}{x}\right) = C \quad .$$

We now have an equation specifying  $y$  as an implicit function of  $x$ . An explicit solution of the differential equation (11.1) is obtained whenever equation (11.15) can be solved for  $y$  in terms of  $x$  and  $C$ .

In summary, the change of variables  $y \rightarrow v$  allows us to transform a differential equation of the form (11.2) to a separable differential equation (11.11) which can be solved by integrating both sides of (11.11) and then solving for  $v$  in terms of  $x$  and then substituting  $\frac{y}{x}$  for  $v$  and solving for  $y$  in terms of  $x$ .

EXAMPLE 11.4.

$$(11.16) \quad y' = \frac{x + y}{x}$$

This equation is homogeneous, since we can re-write it as

$$(11.17) \quad y' = 1 + \frac{y}{x} \quad .$$

We thus take

$$(11.18) \quad F(v) = 1 + v$$

and try to solve

$$(11.19) \quad \frac{dx}{x} = \frac{dv}{F(v) - v} = \frac{dv}{1 + v - v} = dv \quad .$$

Integrating both sides yields

$$(11.20) \quad \ln(x) = v + C = \frac{y}{x} + C$$

or

$$(11.21) \quad y = x(\ln(x) - C) \quad .$$

## 2. Other Substitutions

The substitution method can also be applied in other situations; however, in such cases there usually isn't a clean litmus test that will tell you whether or not a given substitution will help solve the differential equation. Rather one has to resort to trial and error in order to find an appropriate substitution. There is at least a guiding principle though: you want to look for a substitution that will end up simplifying the differential equation. Here are some examples.

EXAMPLE 11.5.

$$(11.22) \quad \frac{dy}{dx} = (x + y + 1)^2 + 3$$

In order to simplify the quadratic expression on the right hand side we'll try the following substitution:

$$z = x + y + 1 \Rightarrow \frac{dz}{dx} = 1 + \frac{dy}{dx} + 0 \Rightarrow \frac{dy}{dx} = \frac{dz}{dx} - 1$$

Substituting  $z$  for  $x + y + 1$  on the right hand side of (11.22) and  $z' - 1$  for  $y'$  on the left hand side of (11.22) we obtain the following equivalent differential equation:

$$\frac{dz}{dx} - 1 = z^2 + 3$$

or

$$\frac{dz}{dx} = z^2 + 4$$

or

$$\frac{dz}{z^2 + 4} = dx$$

This equation is separable, and integrating both sides yields

$$\frac{1}{2} \arctan \frac{1}{2} z = \int \frac{dz}{z^2 + 4} = \int dx + C = x + C$$

or

$$\arctan \left( \frac{z}{2} \right) = 2x + C'$$

or

$$z = 2 \tan (2x + C')$$

or, recalling that  $z = x + y + 1$ ,

$$x + y + 1 = 2 \tan (2x + C')$$

or, solving for  $y$ ,

$$y = 2 \tan (2x + C') - x - 1$$